

Measure	CY	ARHOME Overall	For Comparison			ARHOME FFS/QHP****		By Qualified Health Plan (QHP)**					By Urban/Rural***		By Race/Ethnicity***			
			Mean of Reporting States Medicaid*	NCCA**	AR Medicaid Overall**	Medically Frail (FFS)	QHP	BCBS: Blue Cross Blue Shield	BCBS: Health Advantage	Centene: Ambetter	Centene: QCA	Centene: Qual Choice	Urban	Rural	White	Black	Other	Unknown
Preventive Care																		
Cervical Cancer Screening, 21-64 Years	2019	46.0%	54.1%	60.1%	40.0%	33.9%	47.3%	44.4%	N/A	42.1%	31.0%	30.2%	46.2%	45.9%	45.3%	50.4%	50.9%	41.0%
	2020	43.5%	55.5%	56.8%	41.9%	30.7%	44.6%	41.3%	N/A	38.4%	29.3%	29.6%	43.8%	43.2%	43.0%	48.6%	46.4%	38.0%
	2021	41.7%			43.3%	25.8%	42.7%	42.3%	16.0%	40.0%	30.8%	31.1%	42.0%	41.3%	40.7%	45.9%	46.4%	36.2%
Chlamydia Screening in Women, 21-24 Years	2019	53.9%	59.3%	58.0% Ages 16-24	61.60%	50.2%	54.0%	53.6%	N/A	53.6%	55.5%	55.2%	52.7%	55.5%	49.5%	65.6%	57.0%	50.9%
	2020	52.5%	58.3%	54.5% Ages 16-24	53.7%	54.0%	52.4%	49.7%	N/A	54.7%	52.3%	55.4%	52.4%	52.6%	46.8%	65.0%	50.3%	53.7%
	2021	53.9%			55.5%	53.9%	53.9%	51.3%	50.0%	53.1%	56.8%	57.3%	54.4%	53.1%	48.7%	66.3%	55.2%	51.3%
Breast Cancer Screening, 50-64 Years	2019	50.8%	52.7% Ages 50-64	58.4% Ages 50-74	39.6% Ages 50-74	45.0%	51.6%	54.0%	N/A	49.1%	38.7%	42.2%	50.5%	51.0%	49.0%	55.4%	57.9%	50.7%
	2020	47.7%	53.7% Ages 50-64	53.7% Ages 50-74	42.6% Ages 50-74	38.0%	48.9%	50.9%	N/A	47.1%	40.5%	41.0%	48.2%	47.2%	46.0%	52.8%	52.6%	47.5%
	2021	44.5%			41.7% Ages 50-64	32.9%	45.8%	47.6%	NA	44.4%	39.3%	40.2%	44.7%	44.3%	42.4%	50.7%	47.6%	44.9%
Maternal and Perinatal Care																		
Contraceptive Care – Postpartum Women, Most or Moderately Effective Contraception - 60 Day: 21-44 Years	2019	54.3%	38.4%		38.1%	52.3%	54.4%	54.7%	N/A	53.9%	50.4%	58.4%	52.7%	56.6%	55.3%	54.5%	49.4%	53.6%
	2020	48.9%	40.4%		37.7%	46.2%	49.0%	46.6%	N/A	50.0%	46.5%	49.8%	47.3%	51.3%	52.5%	48.1%	40.7%	43.9%
	2021	45.8%			37.7%	35.2%	45.9%	46.4%	38.7%	44.6%	43.6%	49.8%	42.8%	50.0%	48.2%	44.7%	41.8%	39.6%
Contraceptive Care – All Women, Most or Moderately Effective Contraception: 21-44 Years	2019	25.5%			23.1%	18.3%	26.0%	27.0%	N/A	24.0%	24.3%	24.3%	25.7%	25.3%	25.2%	26.0%	26.4%	25.6%
	2020	23.8%	25.3%		23.6%	17.0%	24.1%	25.2%	N/A	22.3%	22.4%	21.5%	24.1%	23.4%	23.5%	24.2%	23.9%	24.1%
	2021	22.9%			22.8%	14.5%	23.2%	24.6%	19.0%	21.3%	22.4%	22.0%	23.0%	22.7%	22.4%	23.7%	23.3%	23.4%
Low Birth Weight, Percentage of live births weighing < 2,500 grams [‡] (Lower is Better)	2019	10.2%	9.8%††		10.2%			10.2%	N/A	10.5%	9.8%	9.3%	10.3%	10.1%	8.7%	14.8%	8.6%	8.8%
	2020	10.8%			10.6%			11.1%	N/A	10.8%	11.5%	9.6%	11.3%	10.0%	9.7%	15.9%	5.8%	9.7%
	2021																	
Very Low Birth Weight, Percentage of live births weighing < 1,500 grams [‡] (Lower is Better)	2019	1.4%			1.5%			1.2%	N/A	1.7%	1.2%	1.2%	1.3%	1.4%	1.0%	2.4%	0.6%	1.5%
	2020	1.6%			1.3%			1.6%	N/A	1.8%	1.7%	1.2%	1.8%	1.3%	1.3%	2.9%	1.0%	1.0%
	2021																	
Pre-Term Birth, Percentage of live births 17 - 36 weeks gestation [†] (Lower is Better)	2019	13.5%			12.6%			13.3%	N/A	14.0%	14.4%	12.2%	13.3%	13.7%	13.6%	16.1%	10.9%	10.3%
	2020	12.8%			12.4%			13.6%	N/A	14.2%	11.8%	11.2%	13.2%	12.2%	12.8%	15.9%	9.1%	10.6%
	2021																	
Care of Acute and Chronic Conditions																		
Diabetes Short-Term Complications Admission Rate, 19-64 Years (Lower is Better)	2019	26.2	20.6 Ages 18-64		37.3 Ages 18-64	70.4	21.6	14.2	N/A	16.8	16.4	22.4	27.4	24.8	26.6	26.8	20.2	26.7
	2020	21.4	22.2 Ages 18+		20.6 Ages 18-64	44.9	19.3	14.2	N/A	15.5	30.9	27.5	24.0	18.2	22.6	26.2	10.2	17.7
	2021	21.9			21.9 Ages 18-64	58.8	19.3	16.7	23.0	14.6	18.7	17.7	23.0	20.2	22.0	26.4	16.0	18.8
Chronic Obstructive Pulmonary Disease (COPD) or Asthma in Older Adults Admission Rate, 40-64 Years (Lower is Better)	2019	40.9	82.4		121.7	101.2	32.0	24.9	N/A	32.2	18.3	23.4	39.3	42.8	45.8	26.4	33.0	41.1
	2020	23.2	69.4 Ages 40+		33.6	61.6	18.4	14.3	N/A	17.2	19.2	7.7	22.5	24.1	25.6	20.4	8.5	23.4
	2021	19.4			33.3	48.6	16.8	17.5	12.2	17.1	11.7	8.7	15.5	24.1	24.7	14.4	6.8	9.2
Heart Failure Admission Rate, 19-64 Years (Lower is Better)	2019	23.9	31.9 Ages 18-64		47.1 Ages 18-64	88.6	17.8	13.9	N/A	13.5	12.3	13.9	28.1	18.8	19.4	36.8	13.7	28.7
	2020	22.8	31.6 Ages 18+		22.7 Ages 18-64	91.3	17.0	14.4	N/A	16.3	18.3	10.9	27.0	17.4	19.8	36.8	13.8	21.6

Measure	CY	ARHOME Overall	For Comparison			ARHOME FFS/QHP****		By Qualified Health Plan (QHP)**					By Urban/Rural***		By Race/Ethnicity***			
			Mean of Reporting States Medicaid*	NCQA**	AR Medicaid Overall**	Medically Frail (FFS)	QHP	BCBS: Blue Cross Blue Shield	BCBS: Health Advantage	Centene: Ambetter	Centene: QCA	Centene: Qual Choice	Urban	Rural	White	Black	Other	Unknown
	2021	21.7			26.2 Ages 18-64	86.1	17.7	14.8	18.1	18.4	13.1	11.7	25.3	17.1	19.6	34.8	10.8	18.8
Asthma in Younger Adults Admission Rate, 19-39 Years <i>(Lower is Better)</i>	2019	4.8	6.5 Ages 18-39		7.0 Ages 18-39	10.2	4.4	3.1	N/A	3.3	2.1	2.1	5.1	4.5	4.1	9.6	2.4	2.9
	2020	2.1	8.2 Ages 18-39		2.7 Ages 18-39	6.6	1.8	1.6	N/A	2.0	1.7	2.8	2.0	2.2	1.9	4.5	1.4	0.6
	2021	1.7			2.9 Ages 18-39	3.4	1.6	1.8	0.0	1.0	1.2	1.8	1.5	2.0	1.6	3.0	1.1	0.9
Plan All-Cause Readmissions, Observed/Expected Ratio: 19-64 Years <i>(Lower is Better)</i>	2019	0.8506	0.8555 Ages 18-64		0.8906 Ages 18-64	0.9792	0.8194	0.8071	N/A	0.8003	0.7065	0.9174	0.8268	0.8801	0.8635	0.8239	0.7190	0.8502
	2020	0.7743	1.0259 Ages 18-64		1.1297 Ages 18-64	0.9083	0.7435	0.7072	N/A	0.7528	0.4663	0.3911	0.7834	0.7624	0.7967	0.8003	0.7193	0.6705
	2021	0.8457			1.0544 Ages 18-64	1.0546	0.8093	0.7291	7.1528	0.8802	0.9275	0.8545	0.8301	0.8754	0.8318	0.8896	0.7701	0.8879
Asthma Medication Ratio, 19-64 Years	2019	46.9%	55.3%	63.0% Ages 5-64	7.0 Ages 18-39	31.1%	48.5%	48.4%	N/A	45.3%	50.0%	54.5%	50.2%	43.3%	47.6%	47.4%	51.0%	42.9%
	2020	42.4%	53.4%	65.4% Ages 5-64		38.8%	42.7%	43.6%	N/A	36.9%	43.0%	36.1%	45.6%	38.2%	43.3%	40.0%	41.9%	41.7%
	2021	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Behavioral Health Care																		
Initiation of Alcohol and Other Drug Abuse or Dependence Treatment, 19-64 Years	2019	37.9%	41.0% Ages 18-64	44.0% Ages 13+		36.6%	38.2%	37.4%	N/A	38.5%	44.0%	41.5%	37.3%	38.8%	39.1%	31.8%	36.9%	39.5%
	2020	39.2%	43.4% Ages 18+	45.1% Ages 13+	40.7% Ages 18-64	38.5%	39.4%	39.8%	N/A	40.2%	37.4%	38.5%	39.3%	39.2%	40.5%	32.5%	37.7%	41.2%
	2021	40.1%			43.9% Ages 18-64	35.5%	40.6%	41.5%	42.5%	40.8%	38.8%	38.3%	40.4%	39.8%	41.5%	34.9%	39.4%	39.4%
Engagement of Alcohol and Other Drug Abuse or Dependence Treatment, 19-64 Years	2019	8.6%	15.7% Ages 18-64	14.4% Ages 13+		5.2%	9.4%	9.6%	N/A	9.8%	10.3%	8.6%	8.3%	9.0%	9.5%	5.1%	8.6%	8.4%
	2020	9.7%	16.5% Ages 18+	14.1% Ages 13+	7.0% Ages 18-64	6.3%	10.4%	9.5%	N/A	12.0%	9.1%	10.1%	9.2%	10.4%	10.7%	4.6%	9.8%	10.6%
	2021	11.8%			10.2% Ages 18-64	7.2%	12.4%	12.1%	13.2%	13.5%	11.4%	9.8%	12.0%	11.3%	12.8%	7.6%	10.1%	11.8%
Antidepressant Medication Management, Effective Acute Phase Treatment: 19-64 Years	2019	52.9%	51.3% Ages 18-64	55.0% Ages 18+	39.7% Ages 18-64	45.6%	54.7%	55.5%	N/A	56.0%	48.7%	54.8%	52.6%	53.3%	55.0%	40.5%	48.2%	56.6%
	2020	54.0%	52.5% Ages 18+	57.3% Ages 18+	49.5% Ages 18-64	46.5%	55.7%	56.7%	N/A	55.1%	50.8%	52.2%	54.4%	53.4%	56.6%	39.4%	51.9%	56.6%
	2021	58.1%			55.2% Ages 18-64	46.2%	59.7%	59.2%	72.2%	60.7%	57.2%	58.1%	58.8%	57.1%	60.7%	45.4%	56.5%	60.3%
Antidepressant Medication Management, Effective Continuation Phase Treatment: 19-64 Years	2019	37.1%	34.4% Ages 18-64	39.3% Ages 18+	26.1% Ages 18-64	31.3%	38.4%	39.6%	N/A	39.2%	35.6%	35.6%	38.0%	36.0%	39.3%	25.6%	32.0%	39.7%
	2020	38.1%	35.9% Ages 18+	41.6% Ages 18+	33.4% Ages 18-64	30.4%	39.9%	41.3%	N/A	38.3%	35.2%	35.0%	38.2%	38.0%	40.5%	24.6%	37.0%	40.7%
	2021	41.4%			39.4% Ages 18-64	30.4%	42.8%	43.1%	61.1%	42.2%	38.6%	41.9%	41.9%	40.6%	44.0%	27.5%	39.5%	44.6%
Adherence to Antipsychotic Medications for Individuals With Schizophrenia, 19-64 Years	2019	44.1%	61.1%	60.8% Ages 18+	59.4% Ages 18+	43.9%	44.7%	47.2%	N/A	34.8%	65.0%	38.5%	41.1%	47.3%	47.5%	36.6%	41.2%	42.1%
	2020	47.2%	61.2% Ages 18+	61.6% Ages 18+	58.0% Ages 18+	48.1%	46.8%	44.2%	N/A	46.4%	52.1%	43.3%	45.7%	49.1%	50.8%	43.2%	48.5%	39.1%
	2021	41.2%			54.3% Ages 18+	39.9%	42.0%	44.8%	40.0%	39.9%	40.6%	41.0%	38.4%	45.0%	44.8%	33.6%	41.9%	35.4%
Use of Pharmacotherapy for	2019	39.0%			21.8% Ages 18-64	18.9%	43.5%	47.1%	N/A	36.5%	40.2%	45.1%	35.0%	45.4%	42.3%	15.6%	28.6%	34.4%

Measure	CY	ARHOME Overall	For Comparison			ARHOME FFS/QHP****		By Qualified Health Plan (QHP)**					By Urban/Rural***		By Race/Ethnicity***			
			Mean of Reporting States Medicaid*	NCQA**	AR Medicaid Overall**	Medically Frail (FFS)	QHP	BCBS: Blue Cross Blue Shield	BCBS: Health Advantage	Centene: Ambetter	Centene: QCA	Centene: Qual Choice	Urban	Rural	White	Black	Other	Unknown
Use of Pharmacotherapy for Opioid Use Disorder, Overall Total: 19-64 Years	2020	51.3%			47.5% Ages 18-64	32.2%	54.0%	54.0%	N/A	54.1%	55.3%	51.6%	49.4%	54.1%	55.2%	19.9%	45.0%	49.6%
	2021	56.8%			55.6% Ages 18-64	35.4%	58.9%	60.7%	65.4%	57.8%	56.6%	56.1%	54.6%	60.9%	59.8%	28.1%	54.1%	53.9%
Diabetes Screening for People With Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications, 19-64 Years	2019	79.2%	79.9% Ages 18-64	81.7% Ages 18-64		78.0%	80.2%	80.5%	N/A	80.6%	75.2%	81.1%	79.6%	78.8%	80.3%	75.2%	78.9%	78.3%
	2020	77.6%	79.8% Ages 18-64	76.7% Ages 18-64	75.8% Ages 18-64	73.4%	79.0%	78.3%	N/A	79.2%	76.0%	79.4%	77.3%	78.1%	78.1%	79.5%	73.2%	75.7%
	2021	79.7%			80.7% Ages 18-64	76.0%	80.2%	80.2%	78.1%	81.1%	80.5%	79.4%	79.5%	79.8%	80.5%	80.0%	79.9%	74.1%
Use of Opioids at High Dosage in Persons Without Cancer, 19-64 Years <i>(Lower is Better)</i>	2019	1.1%	7.4% Ages 18+		80.3% Ages 18-64	0.5%	1.2%	1.3%	N/A	1.1%	1.1%	0.7%	1.4%	0.7%	1.0%	0.8%	0.7%	1.5%
	2020	1.0%	7.3% Ages 18+		0.7% Ages 18-64	0.4%	1.1%	1.4%	N/A	1.2%	0.3%	0.2%	1.5%	0.6%	1.0%	0.8%	1.5%	1.3%
	2021	0.8%			0.7% Ages 18-64	0.3%	0.8%	0.9%	1.8%	0.9%	0.6%	0.4%	1.1%	0.5%	1.0%	0.4%	0.7%	0.6%
Concurrent Use of Opioids and Benzodiazepines, 19-64 Years <i>(Lower is Better)</i>	2019	20.9%			22.3% Ages 18-64	22.1%	20.6%	21.5%	N/A	17.8%	16.0%	20.0%	21.6%	20.1%	23.7%	11.1%	17.7%	20.8%
	2020	18.9%	15.4% Ages 18+		18.6% Ages 18-64	19.2%	18.8%	20.9%	N/A	16.3%	13.8%	15.0%	19.2%	18.5%	21.2%	11.0%	18.6%	18.3%
	2021	17.2%			17.3% Ages 18-64	15.8%	17.2%	20.1%	15.8%	14.0%	12.3%	11.7%	17.3%	17.0%	19.4%	10.5%	13.8%	16.2%
Follow-Up After Emergency Department Visit for Alcohol and Other Drug Abuse or Dependence, Received Follow-Up Within 30 Days of ED Visit: 19-64 Years	2019	8.7%	20.7% Ages 18-64	19.6% Ages 13+	7.3% Ages 18-64	9.1%	8.6%	8.6%	N/A	11.8%	4.3%	2.9%	8.7%	8.7%	9.1%	7.2%	8.8%	8.7%
	2020	11.0%	22.7% Ages 18+	20.2% Ages 13+	9.7% Ages 18-64	8.4%	11.4%	8.5%	N/A	16.8%	10.3%	9.1%	10.7%	11.5%	12.5%	5.9%	9.0%	11.3%
	2021	11.7%			11.2% Ages 18-64	9.3%	12.0%	8.7%	9.6%	15.9%	12.5%	12.3%	11.5%	12.0%	12.9%	7.1%	13.1%	12.0%
Follow-Up After Emergency Department Visit for Mental Illness, Received Follow-Up Within 30 Days of ED Visit: 19-64 Years	2019	37.3%	54.3% Ages 18-64	55.6% Ages 6+	39.2% Ages 18-64	38.2%	37.3%	41.7%	N/A	35.4%	30.1%	18.6%	33.9%	42.2%	40.3%	26.6%	33.3%	40.5%
	2020	33.0%	54.3% Ages 18+	54.4% Ages 6+	35.2% Ages 18-64	41.3%	30.7%	32.6%	N/A	27.7%	27.8%	33.3%	30.9%	35.9%	35.1%	28.2%	31.0%	31.2%
	2021	28.7%			33.4% Ages 18-64	37.3%	27.2%	26.9%	18.1%	27.7%	26.0%	32.6%	27.3%	30.3%	31.2%	17.3%	38.6%	30.3%
Follow-Up After Hospitalization for Mental Illness: Age 19 to 64, Received Follow-Up Within 30 Days of Discharge: 19-64 Years	2019	37.0%	53.3% Ages 18-64	56.9% Ages 6+	42.0% Ages 18-64	36.2%	38.2%	43.4%	N/A	24.6%	37.2%	35.6%	37.8%	36.0%	36.6%	32.7%	37.4%	41.2%
	2020	36.7%	52.4% Ages 18+	58.9% Ages 6+	37.0% Ages 18-64	37.8%	36.6%	41.6%	N/A	39.0%	23.5%	29.7%	37.2%	36.1%	37.2%	33.1%	38.4%	37.2%
	2021	34.6%			34.8% Ages 18-64	31.9%	35.1%	39.1%	27.9%	31.9%	32.6%	32.5%	34.9%	34.4%	36.3%	30.0%	32.0%	32.2%